No. 9 Providing VA senior managers with a timely and concise summary of important HSR&D findings or publications

May 2003

## PTSD Treatment Important Predictor of Remission in Patients with Substance Use Disorders and PTSD

Approximately one-third of those patients with a substance use disorder (SUD) also have a diagnosis of post-traumatic stress disorder (PTSD). Several programs have been developed to treat comorbid PTSD and SUD, yet little is known about their efficacy. Some programs are based on the theory that addressing PTSD symptoms early in the treatment course prevents SUD relapse. On the other hand, many providers believe that patients must maintain a substantial period of abstinence before PTSD symptoms can be addressed.

Investigators in this study examined whether PTSD treatment received during the first and second year that followed inpatient treatment for substance use disorder would predict 5-year SUD remission. Another focus of the study was to assess 12-Step self-help attendance as an adjunct to predicting 5-year remission. Participants in this study (n = 100) were selected from a larger sample of male veteran inpatients that participated in a multi-site VA program evaluation of SUD treatment. Findings of this study suggest that for patients with both SUD and PTSD, a change in PTSD symptoms may be the most important factor in the remission of both disorders. Patients receiving treatment for PTSD in the first 3 months following discharge from inpatient SUD treatment were more likely to be in remission in year 5 of follow-up, as were patients who received PTSD treatment for a longer period of time during the first year following discharge. In addition, attendance in 12-Step self-help in the first year was positively associated with 5-year remission. These findings indicate that providers focusing on treating PTSD immediately after SUD treatment may improve the chances for long-term remission.

Ouimette P, Moos RH, Finney JW. PTSD treatment and 5-year remission among patients with substance use and posttraumatic stress disorders. Journal of Consulting and Clinical Psychology 2003 Apr;71(2):410-14.

This study was supported by VA's Mental Health Strategic Health Group and HSR&D. Drs. Moos and Finney are part of HSR&D's Center for Health Care Evaluation in Palo Alto, CA.

MANAGEMENT BRIEF is produced by the Management Decision and Research Center, a program within VA's Office of Research & Development, Health Services Research & Development Service. Please take a moment to give us your thoughts and suggestions about Management Brief by sending us an e-mail at MDRC.Boston@med.va.gov.